arXiv:1403.1479v2 [math.CO] 9 Mar 2014

A LOWER BOUND ON THE ENTRIES OF THE PRINCIPAL EIGENVECTOR OF A GRAPH

FELIX GOLDBERG

Abstract. We obtain a lower bound on each entry of the principal eigenvector of a non-regular connected graph.

1. Introduction

The theory of graph spectra, in whose earliest annals we find such illustrious names as Hoffman, Bose, Seidel, and Fiedler, has by now attained a fairly mature stage. Recent expositions of the theory may be found in the books [3, 6, 7, 9].

On the other hand, while the theory of graph eigenvectors may be already out of its infancy, it is still very much in a state of toddlerhood. The purpose of the present note is to make a modest contribution to one of the basic problems of this theory - the description of the entries of the principal eigenvector of a non-regular graph.

2. The problem

Let G be a connected graph on n vertices with adjacency matrix $A \in \mathbb{R}^{n\times n}$. The following facts are widely known (and may be found in each of the references mentioned above):

- A is an irreducible nonnegative matrix.
- The spectral radius $\rho(G)$ of A is a simple eigenvalue.
- The eigenvector $x \in \mathbb{R}^n$ corresponding to ρ is positive entrywise.

We shall refer to $\rho(G)$ as the spectral radius of G and when the context is clear, denote simply $\rho = \rho(G)$. The vector x will be referred to as the *principal eigenvector* of G. An alternative name, which we shall not use here, would be the *Perron vector*.

It is also very well known that:

Date: March 6, 2014.

1991 *Mathematics Subject Classification.* 05C50,15A42,15A18.

Key words and phrases. adjacency matrix, Perron vector, principal eigenvector, induced subgraph.

This research was supported by the Israel Science Foundation (grant number 862/10.)

- If G is regular, then all the entries of x are equal.

Papendieck and Recht in [11] were the first to study the problem of estimating the entries of x in the case that G is non-regular. Before presenting their result, we make an assumption which will be sustained throughout the rest of the note:

Assumption 1. *The vector x is normalized so that $\sum_{i=1}^{n} x_i^2 = 1$.*

Theorem 2. [11] *Let G be a connected graph with principal eigenvector x. Let $x_{\max}$ be the largest entry of x. Then*

$$x_{\max} \leq \frac{1}{\sqrt{2}}.$$

Equality is attained if and only if $G = K_{1,n-1}$ is the star on n vertices.

In fact, Papendieck and Recht's full result is more general, holding for every p-norm ($p \in [1, \infty]$) and depending also on ρ. However, in the case of interest to us, $p = 2$, it reduces to $\frac{1}{\sqrt{2}}$.

3. Known bounds on x

Let us introduce some more notation: denote the degree of the ith vertex of G by d_i. The subgraph of G obtained by deleting the ith vertex (and all edges incident on it) will be denoted as $G_{(i)}$. The spectral radius of $G_{(i)}$ will be denoted by ρ_i. Note that since G is connected, we have by [2, Corollary 2.1.5(b)]:

$$\rho > \rho_i.$$

Cioabă and Gregory [5] have generalized Theorem 2 to give upper bounds on every entry of x.

Theorem 3. [5, Theorem 3.2] *Let G be a connected graph with principal eigenvector x. Then for every $1 \leq i \leq n$:*

$$x_i \leq \frac{1}{\sqrt{1 + \frac{\rho^2}{d_i}}}.$$

Equality is attained if and only if $x_i = x_{\max}$, $d_i = n - 1$, and $G_{(i)}$ is regular.

A natural counterpart to Theorem 3 is given by Li, Wang, and Van Mieghem [8]:

Theorem 4. [8] *Let G be a connected graph with principal eigenvector x. Then for every $1 \leq i \leq n$:*

$$x_i \geq \sqrt{\frac{\rho - \rho_i}{2\rho}}.$$

We remark that additional bounds for $x_{\max}$ and $x_{\min}$ can be found in [5, 10]. There are also in the literature results of a different kind where $\sum_{i \in S} x_i^2$ is estimated from above for subsets $S \subseteq V(G)$ which induce either empty [4, 8] or, more generally, regular subgraphs [1]. When S is a singleton set such bounds reduce to an analogue of Theorem 3.

4. A new lower bound

Our new result is another lower bound on x_i, which is often, but not always, better than Theorem 4:

Theorem 5. *Let G be a connected graph with principal eigenvector x. Then for every $1 \leq i \leq n$:*

$$x_i \geq \frac{1}{\sqrt{1 + \frac{d_i}{(\rho - \rho_i)^2}}}.$$

For the proof we need a lemma:

Lemma 6. [12, p. 148] *Let the Hermitian matrix A be partitioned as*

$$A = \begin{bmatrix} a & b^T \\ b & B \end{bmatrix} \tag{1}$$

and let x be a unit eigenvector of A corresponding to the eigenvalue λ. If λ is not an eigenvalue of B, then

$$|x_1|^2 = \frac{1}{1 + ||(\lambda I - B)^{-1} b||^2}.$$

Proof of Theorem 5. Without loss of generality, let $i = 1$ and suppose that A is partitioned as in (1). Then B is the adjacency matrix of $G_{(1)}$. As observed before: $\rho > \rho_1$. This means that ρ is not an eigenvalue of B and the hypothesis of Lemma 6 is satisfied. Thus we have

$$|x_1|^2 = \frac{1}{1 + ||(\rho I - B)^{-1} b||^2} \geq \frac{1}{1 + ||(\rho I - B)^{-1}||^2 ||b||^2},$$

where $||(\rho I - B)^{-1}||$ is the 2-norm, which is known to be equal to

$$\lambda_{\max}((\rho I - B)^{-1}) = \frac{1}{\lambda_{\min}(\rho I - B)} = \frac{1}{\rho - \lambda_{\max}(B)} = \frac{1}{\rho - \rho_1}.$$

Thus, since $||b||^2 = d_1$ we obtain

$$|x_1|^2 \geq \frac{1}{1 + \frac{d_1}{(\rho - \rho_1)^2}}.$$

□

5. An example

Consider the following graph:



In the table we list the actual values of the principal eigenvector x and the bounds given by all three theorems discussed.

vertex name	vertex degree	Theorem 4	Theorem 5	x_i	Theorem 3
b	6	0.39725	0.45901	0.49917	0.5213
c	6	0.374	0.41636	0.48264	0.5213
g	4	0.29584	0.33114	0.39818	0.44634
a	3	0.18076	0.14959	0.26109	0.39654
e	3	0.25233	0.28276	0.34415	0.39654
i	3	0.18904	0.16325	0.27064	0.39654
d	2	0.17415	0.16949	0.24485	0.33261
f	2	0.13045	0.096049	0.18786	0.33261
h	1	0.044799	0.016093	0.065114	0.24198

As the table makes clear, Theorems 5 and Theorem 4 are, generally speaking, incomparable. Nevertheless, a rule of thumb may be discerned as to when is one better than the other: Theorem 5 works better for vertices of higher degree and Theorem 4 for vertices of low degree. As vertices a, e, i show, however, this rule of thumb is not perfect.

References

[1] M. Andelić and D. M. Cardoso. Spectral characterization of families of split graphs. *Graphs and Combinatorics*, In press.

[2] A. Berman and R. J. Plemmons. *Nonnegative Matrices in the Mathematical Sciences*, volume 9 of *Classics in Applied Mathematics*. SIAM, 1994.

[3] A. E. Brouwer and W. H. Haemers. *Spectra of Graphs*, volume 223 of *Universitext*. Springer, 2012.

[4] S. M. Cioabă. A necessary and sufficient eigenvector condition for a connected graph to be bipartite. *Electron. J. Linear Algebra*, 20:351–353, 2010.

[5] S. M. Cioabă and D. A. Gregory. Principal eigenvectors of irregular graphs. *Electron. J. Linear Algebra*, 16:366–379, 2007.

[6] D. Cvetković, P. Rowlinson, and S. K. Simić. *An Introduction to the Theory of Graph Spectra*, volume 75 of *London Mathematical Society Student Texts*. Cambridge University Press, 2010.

[7] C. Godsil and G. Royle. *Algebraic Graph Theory*, volume 207 of *Graduate Texts in Mathematics*. Springer, 2001.

[8] C. Li, H. Wang, and P. V. Mieghem. Bounds for the spectral radius of a graph when nodes are removed. *Linear Algebra Appl.*, 437(1):319–323, 2012.

[9] P. V. Mieghem. *Graph spectra for complex networks*. Cambridge University Press, 2011.

[10] V. Nikiforov. A spectral condition for odd cycles in graphs. *Linear Algebra Appl.*, 428(7):1492–1498, 2008.

[11] B. Papendieck and P. Recht. On maximal entries in the principal eigenvector of graphs. *Linear Algebra Appl.*, 310(1–3):129–138, 2000.

[12] T. Tao and V. Vu. Random matrices: universality of local eigenvalue statistics. *Acta Math.*, 26(1):127–204, 2011.

Caesarea-Rothschild Institute, University of Haifa, Haifa, Israel
E-mail address: felix.goldberg@gmail.com